arXiv:1211.4433v1 [cs.DM] 19 Nov 2012

An upper bound for the crossing number of bubble-sort graph B_n*

Baigong Zheng, Yuansheng Yang†, Xirong Xu
School of Computer Science and Technology
Dalian University of Technology
Dalian, 116024, P.R. China

Abstract

The *crossing number* of a graph G is the minimum number of pairwise intersections of edges in a drawing of G. Motivated by the recent work [Faria, L., Figueiredo, C.M.H. de, Sýkora, O., Vrt'o, I.: An improved upper bound on the crossing number of the hypercube. J. Graph Theory 59, 145–161 (2008)], we give an upper bound of the crossing number of n-dimensional bubble-sort graph B_n.

Keywords: *Crossing number*; *Drawing*; *Bubble-sort graph*

1 Introduction

Symmetric graphs, such as the n-dimensional Boolean hypercube Q_n, and the cube-connected cycles, have been widely used as processor or communication interconnection networks. For designing, analyzing and improving such networks, Akers and Krishnamurthy proposed a formal group-theoretic model in [1]. Based on the model, the n-dimensional bubble-sort graph is proposed as one of those classes of networks that have better performance, as measured by diameter, connectivity, fault tolerance, etc, than the popular Q_n. Thus, it has drawn a great deal of attention of research (see [2, 4, 9]).

The *n-dimensional bubble-sort graph* B_n has $n!$ vertices labeled by distinct permutations on $\{1, 2, \cdots, n\}$. Let $x = x_1x_2\cdots x_n$ be the vertex of

*The research is supported by NSFC of China (No.60973014, 61170303)
†corresponding author's email : yangys@dlut.edu.cn

B_n. Two vertices $x = x_1x_2\cdots x_n$ and $y = y_1y_2\cdots y_n$ in B_n are adjacent if and only if $x_i = y_{i+1}$ and $x_{i+1} = y_i$ for some i and $x_j = y_j$ for all $j \neq i$ and $i+1$. It is obvious B_n is an $(n-1)$-regular graph.

The crossing number of a graph G, denoted by $cr(G)$, is the smallest number of pairwise crossings of edges among all drawings of G in the plane. In the past thirty years, it turned out that crossing number played an important role not only in various fields of discrete and computational geometry, but also in the design of VLSI circuits [3] and wiring layout problems. However, computing the crossing number was proved to be NP-complete by Garey and Johnson [7]. Thus, it is not surprising that the exact crossing numbers are known for graphs of few families and that the arguments often strongly depend on their structures (see [8, 10]).

Concerned with the upper bound of the crossing number of Q_n, there is a long-standing conjecture proposed by Erdős and Guy [5] in 1973:

$$cr(Q_n) \leq \frac{5}{32}4^n - \lfloor\frac{n^2+1}{2}\rfloor 2^{n-2}.$$

This conjecture remains open till L. Faria, C.M.H de Figueiredo, O. Sýkora, and I. Vrt'o in [6] constructed a good drawing of Q_n which gives the upper bound. However, the upper bound of the crossing number of B_n is still unknown.

Motivated by [6], we give an upper bound of the crossing number of the bubble-sort graph B_n by constructing a drawing of B_n in the plane in this article. In Section 2, we introduce some technical notations and tools, while in Section 3 we give an upper bound of the crossing number of B_n for $n \geq 5$.

2 Preliminaries

We use $V(G)$ and $E(G)$ to denote the vertex set and the edge set of G, respectively. A drawing of G is said to be a *good* drawing, provided that no edge crosses itself, no adjacent edges cross each other, no two edges cross more than once, and no three edges cross in a point. It is well known that the crossing number of a graph is attained only in good drawings of the graph. So we always assume that all drawings throughout this article are good drawings. For a good drawing D of a graph G, we denote by $\nu(D)$ the number of crossings in D. It is clear that $cr(G) \leq \nu(D)$.

Let A and B be two disjoint subsets of an edge set E. The number of the crossings formed by an edge in A and another edge in B is denoted by

$\nu_D(A, B)$ in a drawing D. The number of the crossings that involve a pair of edges in A is denoted by $\nu_D(A)$. Then $\nu_D(A \cup B) = \nu_D(A) + \nu_D(B) + \nu_D(A, B)$.

Next, we present some topological results. Using the results, we derive expressions for counting the crossings of the drawing that is constructed by us in the following section. For $n \geq 6$, let $P = (k_1, k_2, \cdots, k_{n-2})$ be an arbitrary permutation on $\{2, 3, \cdots, n-1\}$. For $0 \leq a \leq n-2$, we define a structure *mesh* $M_{n,a}$ depending on P in the real plane $\mathbb{R}^2$ which is used in the counting-crossing process. Join the points $(0, 1)$, $(0, 2)$, $(0, 3), \cdots, (0, n-1)$ and $(0, n)$ of the vertical real axis with a line. For $1 \leq i \leq a$, let $S_{l,i} = \{l_{ij} : j \in \{1, 2, \cdots, n\} - \{k_i\}\}$ be a group of non-vertical parallel semi-straight lines in the left semi-plane, such that the point $(0, j)$ belongs to l_{ij} and $l_{11}, \cdots, l_{a1}$ lie anticlockwise around point $(0, 1)$. For $a+1 \leq i \leq n-2$, let $S_{r,i} = \{r_{ij} : j \in \{1, 2, \cdots, n\} - \{k_i\}\}$ be a group of non-vertical parallel semi-straight lines in the right semi-plane, such that the point $(0, j)$ belongs to r_{ij}, and $r_{a+1\ 1}, \cdots, r_{n-2\ 1}$ lie clockwise around point $(0, 1)$. The edge $l_{i\ k_i}$ $(1 \leq i \leq a)$ $(r_{i\ k_i}$ $(a+1 \leq i \leq n-2))$ is called the "lost" edge with respect to point $(0, k_i)$. In Figure 2.1 we show drawings of $M_{6,1}$ and $M_{6,2}$ depending on permutation $(2, 4, 5, 3)$.




Figure 2.1: Drawings of $M_{6,1}$ and $M_{6,2}$ depending on $(2, 4, 5, 3)$

By counting the crossings in $M_{n,a}$, we can get

Lemma 2.1. *For positive integers $n \geq 6$ and $0 \leq a \leq n-2$,*

$$\begin{array}{l} \nu_D(S_{l,i_1}, S_{l,i_2}) \\ = \left\{ \begin{array}{l} \binom{n}{2} - (n - k_{i_2}) - (k_{i_1} - 1) = \frac{1}{2}n^2 - \frac{3}{2}n + 1 + k_{i_2} - k_{i_1},\ if\ k_{i_1} < k_{i_2} \\ \binom{n}{2} - (n - k_{i_2}) - (k_{i_1} - 2) = \frac{1}{2}n^2 - \frac{3}{2}n + 2 + k_{i_2} - k_{i_1},\ if\ k_{i_1} > k_{i_2} \end{array} \right. \end{array}$$

for $1 \le i_1 < i_2 \le a$, *and*

$$\begin{array}{rl} & \nu_D(S_{r,i_1}, S_{r,i_2}) \\ = & \begin{cases} \binom{n}{2} - (n-k_{i_2}) - (k_{i_1}-1) = \frac{1}{2}n^2 - \frac{3}{2}n + 1 + k_{i_2} - k_{i_1}, \ if\ k_{i_1} < k_{i_2} \\ \binom{n}{2} - (n-k_{i_2}) - (k_{i_1}-2) = \frac{1}{2}n^2 - \frac{3}{2}n + 2 + k_{i_2} - k_{i_1}, \ if\ k_{i_1} > k_{i_2} \end{cases} \end{array}$$

for $a+1 \le i_1 < i_2 \le n-2$.

For any mesh $M_{n,a}$ depending on $(k_1, k_2, \cdots, k_{n-2})$, let $M'_{n,a}$ be the mesh depending on $(k'_1, k'_2, \cdots, k'_{n-2})$, where $\{k'_1, k'_2, \cdots, k'_a\} = \{k_1, k_2, \cdots, k_a\}$, $\{k'_{a+1}, k'_{a+2}, \cdots, k'_{n-2}\} = \{k_{a+1}, k_{a+2}, \cdots, k_{n-2}\}$ and $k'_i < k'_j$ for $1 \le i < j \le a$ and $a+1 \le i < j \le n-2$ separately. Then we have the following lemmas.

Lemma 2.2. *For positive integers* $n \ge 6$ *and* $0 \le a \le n-2$, $\nu_D(E(M_{n,a})) \le \nu_D(E(M'_{n,a}))$.

Proof. Let $\beta = \beta_l + \beta_r$, where β_l (or β_r) is the number of inversions in permutation section $(k_1, k_2, \cdots, k_a)$ (or $(k_{a+1}, k_{a+2}, \cdots, k_{n-2})$). Let $\beta' = \beta'_l + \beta'_r$, where β'_l (or β'_r) is the number of inversions in permutation section $(k'_1, k'_2, \cdots, k'_a)$ (or $(k'_{a+1}, k'_{a+2}, \cdots, k'_{n-2})$). Then $\beta' = 0$. If $\beta = 0$, then $(k_1, k_2, \cdots, k_{n-2}) = (k'_1, k'_2, \cdots, k'_{n-2})$, and $\nu_D(E(M_{n,a})) = \nu_D(E(M'_{n,a}))$.

Now we consider the case for $\beta = m > 0$. Let $P^0 = (k^0_1, k^0_2, \cdots, k^0_{n-2}) = (k_1, k_2, \cdots, k_{n-2})$. For $1 \le t \le m$, suppose $(k^{t-1}_i, k^{t-1}_{i+1})$ be the first adjacent inversion in permutation section $(k^{t-1}_1, k^{t-1}_2, \cdots, k^{t-1}_a)$ or $(k^{t-1}_{a+1}, k^{t-1}_{a+2}, \cdots, k^{t-1}_{n-2})$, let $P^t = (k^t_1, k^t_2, \cdots, k^t_{n-2})$, where $k^t_i = k^{t-1}_{i+1}$, $k^t_{i+1} = k^{t-1}_i$ and $k^t_j = k^{t-1}_j$ for $1 \le j \le i-1$ and $i+2 \le j \le n-2$.

For $0 \le t \le m$, Let $\beta^t = \beta^t_l + \beta^t_r$, where β^t_l (or β^t_r) is the number of inversions in permutation section $(k^t_1, k^t_2, \cdots, k^t_a)$ (or $(k^t_{a+1}, k^t_{a+2}, \cdots, k^t_{n-2})$). Then $\beta^0 = \beta = m$, $\beta^t = \beta^{t-1} - 1$ and $\beta^m = 0 = \beta'$.

Let $M^t_{n,a}$ be the mesh depending on P^t. Then $P^m = (k^m_1, k^m_2, \cdots, k^m_{n-2})$ $= (k'_1, k'_2, \cdots, k'_{n-2})$, $M^0_{n,a} = M_{n,a}$ and $M^m_{n,a} = M'_{n,a}$. For $1 \le i \le a$, let $S^t_{l,i} = \{l_{ij} : j \in \{1, 2, \cdots, n\} - \{k^t_i\}\}$, and for $a+1 \le i \le n-2$ let $S^t_{r,i} = \{r_{ij} : j \in \{1, 2, \cdots, n\} - \{k^t_i\}\}$.

By Lemma 2.1,

$$\begin{array}{l} \nu_D(E(M^t_{n,a})) - \nu_D(E(M^{t-1}_{n,a})) \\ = \begin{cases} \nu_D(S^t_{l,i}, S^t_{l,i+1}) - \nu_D(S^{t-1}_{l,i}, S^{t-1}_{l,i+1}), & if\ 1 \le i \le a-1 \\ \nu_D(S^t_{r,i}, S^t_{r,i+1}) - \nu_D(S^{t-1}_{r,i}, S^{t-1}_{r,i+1}), & if\ a+1 \le i \le n-3 \end{cases} \\ = \left(\frac{1}{2}n^2 - \frac{3}{2}n + 1 + k^t_{i+1} - k^t_i\right) - \left(\frac{1}{2}n^2 - \frac{3}{2}n + 2 + k^{t-1}_{i+1} - k^{t-1}_i\right) \\ = k^t_{i+1} - k^t_i - k^{t-1}_{i+1} + k^{t-1}_i - 1 \\ = 2k^{t-1}_i - 2k^{t-1}_{i+1} - 1 > 0. \end{array}$$

Hence $\nu_D(E(M_{n,a}^{t-1})) < \nu_D(E(M_{n,a}^{t}))$. It follows $\nu_D(E(M_{n,a})) = \nu_D(E(M_{n,a}^{0}))$ $< \nu_D(E(M_{n,a}^{1})) < \cdots < \nu_D(E(M_{n,a}^{m})) = \nu_D(E(M'_{n,a}))$. □

Lemma 2.3. *For integers $m \geq 4$ and $a \geq 1$,*

$$\begin{aligned}
&\nu_D(E(M_{n,a}))\\
&\leq \begin{cases}
\frac{1}{8}n^4 - \frac{25}{24}n^3 + 3n^2 - \frac{23}{6}n + 2, & if \quad n = 2m,\ a = m-1,\\
\frac{1}{8}n^4 - \frac{25}{24}n^3 + \frac{7}{2}n^2 - \frac{29}{6}n + 2, & if \quad n = 2m,\ a \in \{m-2, m\},\\
\frac{1}{8}n^4 - \frac{25}{24}n^3 + \frac{25}{8}n^2 - \frac{95}{24}n + \frac{7}{4}, & if \quad n = 2m-1,\ a \in \{m-2, m-1\}.
\end{cases}
\end{aligned}$$

Proof. Firstly, we give a proper permutation, called $(k_1'', k_2'', \cdots, k_{n-2}'')$, depending on which the mesh structure is denoted by $M_{n,a}''$.

$$\begin{aligned}
&(k_1'', k_2'', k_3'', \cdots, k_a'')(k_{a+1}'', k_{a+2}'', k_{a+3}'', \cdots, k_{n-2}'')\\
= &\begin{cases}
(2,4,6,\cdots,n-2)(3,5,7,\cdots,n-1), & if \quad n = 2m,\ a = m-1,\\
(2,3,5,7,\cdots,n-1)(4,6,8,\cdots,n-2), & if \quad n = 2m,\ a = m,\\
(4,6,8,\cdots,n-2)(2,3,5,7,\cdots,n-1), & if \quad n = 2m,\ a = m-2,\\
(2,4,6,\cdots,n-1)(3,5,7,\cdots,n-2), & if \quad n = 2m-1,\ a = m-1,\\
(3,5,7,\cdots,n-2)(2,4,6,\cdots,n-1), & if \quad n = 2m-1,\ a = m-2.
\end{cases}
\end{aligned}$$

Then the permutation $(k_1'', k_2'', \cdots, k_{n-2}'')$ satisfies $k_i'' < k_j''$ for $1 \leq i < j \leq a$ and $a+1 \leq i < j \leq n-2$ separately.

We will prove Lemma 2.3 holds for $(n, a) = (2m, m)$. By a similar argument, we can prove Lemma 2.3 holds for the other cases of (n, a) and we leave it to readers.

By Lemmas 2.1 and 2.2,

$$\begin{aligned}
&\nu_D(E(M_{n,a}))\\
&\leq \nu_D(E(M'_{n,a}))\\
&= \sum_{1\leq i<j\leq a} \nu_D(S'_{l,i}, S'_{l,j}) + \sum_{a+1\leq i<j\leq n-2} \nu_D(S'_{r,i}, S'_{r,j})\\
&= \left(\tfrac{1}{2}n^2 - \tfrac{3}{2}n + 1\right)\left[\binom{a}{2} + \binom{n-2-a}{2}\right] + \sum_{1\leq i<j\leq a}(k'_j - k'_i) + \sum_{a+1\leq i<j\leq n-2}(k'_j - k'_i)\\
&= \tfrac{1}{8}(n^4 - 9n^3 + 32n^2 - 48n + 24) + \sum_{i=1}^{\frac{n}{2}}(2i - 1 - \tfrac{n}{2})k'_i + \sum_{i=1}^{\frac{n}{2}-2}(2i + 1 - \tfrac{n}{2})k'_{i+a}\\
&= \tfrac{1}{8}(n^4 - 9n^3 + 32n^2 - 48n + 24) + \sum_{i=1}^{\frac{n}{2}}(2i - 1)k'_i + \sum_{i=1}^{\frac{n}{2}-2}(2i + 1)k'_{i+a} - \sum_{i=1}^{n-2}\tfrac{n}{2}k'_i\\
&= \tfrac{1}{8}(n^4 - 11n^3 + 34n^2 - 44n + 24) + \sum_{i=1}^{\frac{n}{2}}(2i - 1)k'_i + \sum_{i=1}^{\frac{n}{2}-2}(2i + 1)k'_{i+a}.
\end{aligned}$$

Now we will prove $\sum_{i=1}^{\frac{n}{2}}(2i-1)k'_i + \sum_{i=1}^{\frac{n}{2}-2}(2i+1)k'_{i+a} \leq \sum_{i=1}^{\frac{n}{2}}(2i-1)k''_i + \sum_{i=1}^{\frac{n}{2}-2}(2i+1)k''_{i+a}$. We define

$$\theta(i) = \begin{cases} 2i-1 & if \ \ 1 \leq i \leq a, \\ 2(i-a)+1 & if \ \ a+1 \leq i \leq n-2. \end{cases}$$

Let $P_0 = (k_1^0, k_2^0, \cdots, k_{n-2}^0) = (k'_1, k'_2, \cdots, k'_{n-2})$. For $1 \leq t \leq n-3$, suppose $k_{i_t}^{t-1} = n-t$, let $j_t = \frac{n}{2} - \frac{t-1}{2}$ for odd t, $j_t = n - \frac{t}{2} - 1$ for even t and $P_t = (k_1^t, k_2^t, \cdots, k_{n-2}^t)$ where $k_{j_t}^t = k_{i_t}^{t-1}$, $k_{i_t}^t = k_{j_t}^{t-1}$ and $k_s^t = k_s^{t-1}$ for $1 \leq s \leq n-2, s \neq i_t$ and $s \neq j_t$. Then, for even t

$$\begin{aligned} &(k_{\frac{n}{2}-\frac{t}{2}+1}^{t-1}, k_{\frac{n}{2}-\frac{t}{2}+2}^{t-1}, \cdots, k_{\frac{n}{2}}^{t-1})(k_{n-\frac{t}{2}}^{t-1}, k_{n-\frac{t}{2}+1}^{t-1}, \cdots, k_{n-2}^{t-1}) \\ =&(n-t+1, n-t+3, \cdots, n-1)(n-t+2, n-t+4, \cdots, n-2), \quad (2.1) \end{aligned}$$

and for odd t

$$\begin{aligned} &(k_{\frac{n}{2}-\frac{t-1}{2}+1}^{t-1}, k_{\frac{n}{2}-\frac{t-1}{2}+2}^{t-1}, \cdots, k_{\frac{n}{2}}^{t-1})(k_{n-\frac{t+1}{2}}^{t-1}, k_{n-\frac{t+1}{2}+1}^{t-1}, \cdots, k_{n-2}^{t-1}) \\ =&(n-t+2, n-t+4, \cdots, n-1)(n-t+1, n-t+3, \cdots, n-2). \quad (2.2) \end{aligned}$$

Furthermore, we have $k_{i_t}^t = k_{j_t}^{t-1} \leq n-t = k_{i_t}^{t-1} = k_{j_t}^t$ and $P_{n-3} = (k''_1, k''_2, \cdots, k''_{n-2})$.

There are two cases depending on t.

Case 1. t is even. Then $t-1$ is odd. By expression (1), $1 \leq i_t \leq \frac{n}{2} - \frac{t}{2} + 1 - 1 = \frac{n-t}{2}$ or $a+1 \leq i_t \leq n - \frac{t}{2} - 1$. Hence,

$$\begin{aligned} \theta(i_t) &\leq \begin{cases} 2 \cdot \frac{n-t}{2} - 1, & if \ \ 1 \leq i_t \leq \frac{n-t}{2} \\ 2 \cdot (n - \frac{t}{2} - 1 - a) + 1, & if \ \ a+1 \leq i_t \leq n - \frac{t}{2} - 1 \end{cases} \\ &= n-t-1 \\ &= 2(n - \frac{t}{2} - 1 - a) + 1 \\ &= \theta(j_t) \end{aligned}$$

Case 2. t is odd. Then $t-1$ is even. By expression (2), $1 \leq i_t \leq \frac{n}{2} - \frac{t-1}{2} + 1 - 1 = \frac{n}{2} - \frac{t-1}{2}$ or $a+1 \leq i_t \leq n - \frac{t+1}{2} - 1 = n - \frac{t+3}{2}$. Hence,

$$\begin{aligned} \theta(i_t) &\leq \begin{cases} 2 \cdot (\frac{n}{2} - \frac{t-1}{2}) - 1, & if \ \ 1 \leq i_t \leq \frac{n}{2} - \frac{t-1}{2} \\ 2 \cdot (n - \frac{t+3}{2} - a) + 1, & if \ \ a+1 \leq i_t \leq n - \frac{t+3}{2} \end{cases} \\ &\leq n-t \\ &= 2(\frac{n}{2} - \frac{t-1}{2}) - 1 \\ &= \theta(j_t) \end{aligned}$$

So $\theta(i_t) \leq \theta(j_t)$ for all $1 \leq t \leq n-3$.

For $0 \leq t \leq n-3$, let $f(P_t) = \sum_{i=1}^{\frac{n}{2}}(2i-1)k_i^t + \sum_{i=1}^{\frac{n}{2}-2}(2i+1)k_{i+a}^t$. Then for $1 \leq t \leq n-3$,

$$\begin{aligned} f(P_t) - f(P_{t-1}) &= \theta(i_t)k_{i_t}^t + \theta(j_t)k_{j_t}^t - (\theta(i_t)k_{i_t}^{t-1} + \theta(j_t)k_{j_t}^{t-1}) \\ &= \theta(i_t)(k_{i_t}^t - k_{i_t}^{t-1}) + \theta(j_t)(k_{j_t}^t - k_{j_t}^{t-1}) \\ &= (\theta(i_t) - \theta(j_t))(k_{i_t}^t - k_{j_t}^t) \geq 0. \end{aligned}$$

So there is $f(P_{t-1}) \leq f(P_t)$. It follows $f(P_0) \leq f(P_1) \leq \cdots \leq f(P_{n-3})$. Hence,

$$\begin{aligned} &\nu_D(E(M_{n,a})) \\ &\leq \tfrac{1}{8}(n^4 - 11n^3 + 34n^2 - 44n + 24) + \sum_{i=1}^{\frac{n}{2}}(2i-1)k_i' + \sum_{i=1}^{\frac{n}{2}-2}(2i+1)k_{i+a}' \\ &= \tfrac{1}{8}(n^4 - 11n^3 + 34n^2 - 44n + 24) + f(P_0) \\ &\leq \tfrac{1}{8}(n^4 - 11n^3 + 34n^2 - 44n + 24) + f(P_{n-3}) \\ &= \tfrac{1}{8}(n^4 - 11n^3 + 34n^2 - 44n + 24) + \sum_{i=1}^{\frac{n}{2}}(2i-1)k_i'' + \sum_{i=1}^{\frac{n}{2}-2}(2i+1)k_{i+a}'' \\ &= \tfrac{1}{8}n^4 - \tfrac{25}{24}n^3 + \tfrac{7}{2}n^2 - \tfrac{29}{6}n + 2. \end{aligned}$$

□

3 Upper bound for $cr(B_n)$

In Figure 3.1, 3.2 and 3.3, we show a drawing D_2 (D_3, D_4, D_5, D_6) of B_2 (B_3, B_4, B_5, B_6) containing 0 (0, 0, 120, 5196) crossings. So we can obtain the following theorem:




Figure 3.1: Some drawings of B_2, B_3 and B_4




Figure 3.2: A drawing of B_5


Theorem 3.1. *$cr(B_2) = cr(B_3) = cr(B_4) = 0$, $cr(B_5) \leq 120$ and $cr(B_6) \leq 5196$.*

Our main result is Theorem 3.2.

Theorem 3.2. *For $n \geq 7$,*

$$cr(B_n) \leq ((n-1)!)^2 \left[\frac{127}{300} + \frac{5}{24(n-4)!} + \sum_{i=3}^{n-5} \frac{1}{3i!} + \sum_{i=2j,j\geq 4}^{n} \left(\frac{1}{8(i-3)!} - \frac{1}{4(i-2)!} + \frac{1}{4(i-1)!(i-1)} \right) \right].$$

To prove Theorem 3.2, we first need some definitions and observations.



$\nu(D_6) \leq 5196$

Figure 3.3: A drawing of B_6

Definition 3.1. *For any vertex $v \in V(B_n)$, we denote v^1, $v^2, \cdots$, v^{n+1} as $n+1$ vertices of B_{n+1} satisfying $v^i = v_1 v_2 \cdots v_{i-1}(n+1)v_i \cdots v_n$ $(1 \leq i \leq n+1)$.*

Observation 3.1. *For any vertex $v \in V(B_n)$ and $v^i \in V(B_{n+1})$ $(1 \leq i \leq n+1)$, there are n edges $\{v^1v^2, v^2v^3, \cdots, v^nv^{n+1}\}$ resulting in a path with $n+1$ vertices.*

Definition 3.2. *For $n \geq 6$, let B'_n be a subgraph of B_n satisfying :* (1) *$V(B'_n)$ is a subset of $V(B_n)$ and $V(B'_n)$=$\{v$: the label of v contains $(1,2,3,4), (1,2,4,3), (1,4,2,3)$ or $(4,1,2,3)$ as a subpermutation$\}$* (2) *$uv \in E(B'_n)$, **iff** $uv \in E(B_n)$ and at least one of u and v is in $V(B'_n)$. A drawing of B'_n*

is denoted by D'_n.

Let D_n be a drawing of B_n containing six D'_n, and we have

Observation 3.2. *By symmetry* (*see Figure* 3.3 *and* 3.4), $\nu(D_n) = 6 * \nu(D'_n)$.

In Figure 3.4, we show a drawing D'_6. Based on the drawing D'_6, we construct 5 real number axes R_i $(1 \leq i \leq 5)$ as shown in Figure 3.4. All of the axes are parallel and all the vertices of B'_n are drawn precisely on R_i.




Figure 3.4: A drawing D'_6

Definition 3.3. *Let x be a vertex of B'_n on axis R_i and $e = xy$ an edge of B'_n in D'_n.* (1) *If the edge e shots away the axis R_i left from the vertex x, i.e. the part of e in a small enough neighborhood of x lies in the left of the axis R_i, we call xy an **'l-arc'** with respect to x.* (2) *If e shots away the axis R_i right from the vertex x, we call xy an **'r-arc'** with respect to x.* (3) *We write $l(x) = |\{xy \in E(B'_n) : xy \text{ is an l-arc with respect to } x\}|$ and $r(x) = |\{xy \in E(B'_n) : xy \text{ is an r-arc with respect to } x\}|$.*

Next we will construct D'_{n+1} from D'_n ($n \geq 6$) and compute the crossing number $\nu(D'_{n+1})$.

For $1 \leq i \leq 5$, let v be the j-th vertex lying on R_i (from the top down). We replace v of B'_n in the "small" neighborhood of v in the drawing D'_n by path $P_{v^{n+1}v^n\ldots v^1}$ ($P_{v^1v^2\ldots v^{n+1}}$) on R_i for odd j (even j) (See Figure 3.4). Now every drawn edge e in D'_n which started in v will be replaced by n "parallel" edges (bunch) and drawn along the original edge e. Notice that, in this case, locally we have a drawing of mesh. Doing this carefully we get a drawing D'_{n+1} for $n \geq 6$.

To construct D'_{n+1} from D'_n, we use three kinds of mesh structures defined as follows:

Definition 3.4. *A mesh $M_{n,a}$ ($0 \leq a \leq n-2$) is called a **11-structure** mesh, if for $1 \leq j \leq \lfloor \frac{n}{2} \rfloor$, all edges $v^{2j-1}v^{2j}$ are drawn on the left of R_i, and all edges $v^{2j}v^{2j+1}$ are drawn on the right of R_i ($1 \leq i \leq 5$) (See Figure 3.5).*





Figure 3.5: An example of 11-structure mesh for $n = 6$

Definition 3.5. *A mesh $M_{n,a}$ ($0 \leq a \leq n-2$) is called a **20-structure** (**02-structure**) mesh, if n is even and for $1 \leq j \leq n-1$, all edges v^jv^{j+1} are drawn on the left (right) of R_i ($1 \leq i \leq 5$) (See Figure 3.6).*

Now we introduce a *balanced* drawing combined with 11-structure, 20-structure and 02-structure meshes. For $n \geq 6$, every vertex $v \in V(B'_n)$ will be replaced by the path $P_{v^{n+1}v^n\ldots v^1}$ (or $P_{v^1v^2\ldots v^{n+1}}$) of a specific mesh



20-structure 02-structure

Figure 3.6: Examples of 20-structure mesh and 02-structure mesh for $n = 6$

$M_{n+1,l(v)}(v)$ depending on $l(v) - r(v)$ as shown in Table 3.1. We assume that all drawings D'_n $(n \geq 6)$ in the following part are balanced drawings. A part of D'_8 containing 11-structure, 20-structure and 02-structure meshes is shown in Figure 3.7.

Table 3.1: Every vertex in B'_n is replaced by the $(n+1)$-path of a specific mesh

n	$l(v) - r(v)$	$specific\ mesh$
even	$-1, 1$	$11 - structure$
odd	0	$11 - structure$
odd	-2	$20 - structure$
odd	2	$02 - structure$

Lemma 3.1. *For any vertex v in the drawing D'_n with $n \geq 6$, $l(v) - r(v) \in \{0, 2, -2\}$ for odd n, and $l(v) - r(v) \in \{1, -1\}$ for even n.*

Proof. We prove Lemma 3.1 by induction on n.

1) For $n = 6$, there is $l(v) - r(v) \in \{1, -1\}$ for any vertex v in the drawing D'_n (See Figure 3.4). Hence Lemma 3.1 holds for $n = 6$.

2) Assume Lemma 3.1 holds for k. Now consider the case for $k + 1$. There are two cases depending on whether k is even or odd.

Case 1. k is even. v is replaced by a 11-structure mesh.

For $2 \leq i \leq k$, considering the lost edge with respect to v^i, we have $l(v) - 1 + 1 \leq l(v^i) \leq l(v) + 1$ and $r(v) - 1 + 1 \leq r(v^i) \leq r(v) + 1$. For $i = 1$ or $k + 1$, since there is no lost edge with respect to v^i, we have $l(v) \leq l(v^i) \leq l(v)+1$ and $r(v) \leq r(v^i) \leq r(v)+1$. It follows, $|l(v^i)-r(v^i)| \leq$




Figure 3.7: A part of D'_8 containing 11-structure, 20-structure and 02-structure meshes

$|l(v) - r(v)| + 1 = 2$ for all $1 \leq i \leq k+1$. Since $l(v^i) + r(v^i) = k$ is even, we have $l(v^i) - r(v^i) \in \{0, -2, 2\}$.

Case 2. k is odd.

Case 2.1. $l(v) - r(v) = 0$. v is replaced by a 11-structure mesh.

For $2 \leq i \leq k$, we have $l(v) - 1 + 1 \leq l(v^i) \leq l(v) + 1$ and $r(v) - 1 + 1 \leq r(v^i) \leq r(v) + 1$. For $i = 1$ or $k+1$, we have $l(v) \leq l(v^i) \leq l(v) + 1$ and $r(v) \leq r(v^i) \leq r(v) + 1$. It follows $|l(v^i) - r(v^i)| \leq |l(v) - r(v)| + 1 = 1$ for all $1 \leq i \leq k+1$.

Case 2.2. $l(v) - r(v) = 2$. v is replaced by a 02-structure mesh.

For $2 \leq i \leq k$, we have $l(v) - 1 \leq l(v^i) \leq l(v)$ and $r(v) - 1 + 2 \leq r(v^i) \leq r(v) + 2$. It follows $-1 = l(v) - 1 - r(v) - 2 \leq l(v^i) - r(v^i) \leq l(v) - r(v) - 1 = 1$. For $i = 1$ or $k+1$, we have $l(v^i) = l(v)$ and $r(v^i) = r(v) + 1$. Then $l(v^i) - r(v^i) = l(v) - r(v) - 1 = 1$. It follows $|l(v^i) - r(v^i)| \leq 1$ for all $1 \leq i \leq k+1$.

Case 2.3. $l(v) - r(v) = -2$. v is replaced by a 20-structure mesh.

For $2 \leq i \leq k$, we have $l(v) - 1 + 2 \leq l(v^i) \leq l(v) + 2$ and $r(v) - 1 \leq r(v^i) \leq r(v)$. It follows $-1 = l(v) + 1 - r(v) \leq l(v^i) - r(v^i) \leq l(v) + 2 - r(v) + 1 = 1$. For $i = 1$ or $k+1$, we have $l(v^i) = l(v) + 1$ and $r(v^i) = r(v)$. Then $l(v^i) - r(v^i) = l(v) + 1 - r(v) = -1$. It follows $|l(v^i) - r(v^i)| \leq 1$ for all $1 \leq i \leq k+1$.

By Cases 2.1-2.3, we have $|l(v^i) - r(v^i)| \leq 1$ for odd k. Since $l(v^i) + r(v^i) = k$ is odd, we have $l(v^i) - r(v^i) \in \{-1, 1\}$.

By 1) and 2), Lemma 3.1 holds for all $n \geq 6$. □

Lemma 3.2. *For even $n \geq 6$ and any vertex v in the drawing D'_n, there is $|\{v^i : l(v^i) - r(v^i) \in \{2, -2\}\}| = \frac{n}{2}$ and $|\{v^i : l(v^i) - r(v^i) = 0\}| = \frac{n}{2} + 1$ in D'_{n+1}.*

Proof. By Lemma 3.1, $l(v) - r(v) \in \{-1, 1\}$ for even $n \geq 6$.

Case 1. $(l(v), r(v)) = (\frac{n}{2} - 1, \frac{n}{2})$.

$$
\begin{array}{ll}
 & |\{v^i : l(v^i) - r(v^i) \in \{2, -2\}\}| \\
= & |\{v^i : l(v^i) - r(v^i) = -2\}| \\
= & |\{v^i : (l(v^i), r(v^i)) = (\frac{n}{2} - 1 - 1 + 1, \frac{n}{2} + 1)\} \cup \\
 & \{v^{n+1} : (l(v^{n+1}), r(v^{n+1})) = (\frac{n}{2} - 1, \frac{n}{2} + 1)\}| \\
= & |\{v^i : (l(v^i), r(v^i)) = (\frac{n}{2} - 1, \frac{n}{2} + 1)\}| + \\
 & |\{v^{n+1} : (l(v^{n+1}), r(v^{n+1})) = (\frac{n}{2} - 1, \frac{n}{2} + 1)\}| \\
= & l(v) + 1 = \frac{n}{2} - 1 + 1 = \frac{n}{2}.
\end{array}
$$

$$
\begin{array}{ll}
 & |\{v^i : l(v^i) - r(v^i) = 0\}| \\
= & |\{v^i : (l(v^i), r(v^i)) = (\frac{n}{2} - 1 + 1, \frac{n}{2} - 1 + 1)\} \cup \\
 & \{v^1 : (l(v^1), r(v^1)) = (\frac{n}{2} - 1 + 1, \frac{n}{2})\}| \\
= & |\{v^i : (l(v^i), r(v^i)) = (\frac{n}{2}, \frac{n}{2})\}| + |\{v^1 : (l(v^1), r(v^1)) = (\frac{n}{2}, \frac{n}{2})\}| \\
= & r(v) + 1 = \frac{n}{2} + 1.
\end{array}
$$

Case 2. $(l(v), r(v)) = (\frac{n}{2}, \frac{n}{2} - 1)$.

$$
\begin{array}{ll}
 & |\{v^i : l(v^i) - r(v^i) = 0\}| \\
= & |\{v^i : (l(v^i), r(v^i)) = (\frac{n}{2} - 1 + 1, \frac{n}{2} - 1 + 1)\} \cup \\
 & \{v^{n+1} : (l(v^{n+1}), r(v^{n+1})) = (\frac{n}{2}, \frac{n}{2} - 1 + 1)\}| \\
= & |\{v^i : (l(v^i), r(v^i)) = (\frac{n}{2}, \frac{n}{2})\}| + |\{v^{n+1} : (l(v^{n+1}), r(v^{n+1})) = (\frac{n}{2}, \frac{n}{2})\}| \\
= & l(v) + 1 = \frac{n}{2} + 1.
\end{array}
$$

$$
\begin{array}{ll}
 & |\{v^i : l(v^i) - r(v^i) \in \{2, -2\}\}| \\
= & |\{v^i : l(v^i) - r(v^i) = 2\}| \\
= & |\{v^i : (l(v^i), r(v^i)) = (\frac{n}{2} + 1, \frac{n}{2} - 1 - 1 + 1)\} \cup \\
 & \{v^1 : (l(v^1), r(v^1)) = (\frac{n}{2} + 1, \frac{n}{2} - 1)\}| \\
= & |\{v^i : (l(v^i), r(v^i)) = (\frac{n}{2} + 1, \frac{n}{2} - 1)\}| + \\
 & |\{v^1 : (l(v^1), r(v^1)) = (\frac{n}{2} + 1, \frac{n}{2} - 1)\}| \\
= & r(v) + 1 = \frac{n}{2} - 1 + 1 = \frac{n}{2}.
\end{array}
$$

□

To construct D'_{n+1} from D'_n, it can be obtained that there are totally 2 types of crossings in D'_{n+1}.

Type 1. Crossings produced by the crossings in D'_n. As shown in Figure 3.8, each crossing in D'_n will be replaced by n^2 crossings in D'_{n+1}.




Figure 3.8: An example of the crossings of Type 1 for $n = 6$

Type 2. Crossings produced by replacing a vertex of D'_n by $n+1$ vertices in D'_{n+1} (See Figure 2.1).

Then $\nu(D'_{n+1})$ will be $n^2\nu(D'_n)$ plus the sum of the numbers of the crossings in the small neighborhood of the $M_{n+1,l(v)}(v)$ for all vertices $v \in V(B'_n)$.

Lemma 3.3. *For $n = 2m - 2$ $(m \geq 4)$,*

$$\nu(D'_{n+1}) \leq n^2\nu(D'_n) + \frac{n!}{144}(3n^4 - 13n^3 + 18n^2 - 8n).$$

Proof. By Lemma 3.1, $l(v) - r(v) \in \{-1, 1\}$ for any vertex v in D'_n. It follows $l(v) \in \{m-2, m-1\}$, by Lemma 2.3, we have

$$\begin{aligned}
&\nu(D'_{n+1}) \\
&\leq n^2\nu(D'_n) + \tfrac{n!}{6} \cdot \left[\tfrac{1}{8}(n+1)^4 - \tfrac{25}{24}(n+1)^3 + \tfrac{25}{8}(n+1)^2 - \tfrac{95}{24}(n+1) + \tfrac{7}{4}\right] \\
&= n^2\nu(D'_n) + \tfrac{n!}{144}(3n^4 - 13n^3 + 18n^2 - 8n)
\end{aligned}$$

for $n = 2m - 2$ and $m \geq 4$. □

Lemma 3.4. *For $n = 2m - 1$ $(m \geq 4)$,*

$$\nu(D'_{n+1}) \leq n^2\nu(D'_n) + \frac{(n-1)!}{144}(3n^5 - 13n^4 + 21n^3 - 17n^2 + 6).$$

Proof. By Lemma 3.1, $l(v) - r(v) \in \{0, -2, 2\}$ for any vertex v in D'_n. It follows $l(v) \in \{m, m-2, m-1\}$, by Lemmas 2.3 and 3.2, we have

$$\begin{aligned}
&\nu(D'_{n+1}) \\
&\leq n^2\nu(D'_n) + \tfrac{n!}{6} \cdot \tfrac{1}{n} \left\{ \tfrac{n+1}{2} \left[\tfrac{1}{8}(n+1)^4 - \tfrac{25}{24}(n+1)^3 + 3(n+1)^2 - \tfrac{23}{6}(n+1)\right.\right. \\
&\quad \left. + 2\right] + \tfrac{n-1}{2} \left[\tfrac{1}{8}(n+1)^4 - \tfrac{25}{24}(n+1)^3 + \tfrac{7}{2}(n+1)^2 - \tfrac{29}{6}(n+1) + 2\right] \Big\} \\
&= n^2\nu(D'_n) + \tfrac{(n-1)!}{144}(3n^5 - 13n^4 + 21n^3 - 17n^2 + 6)
\end{aligned}$$

for $n = 2m - 1$ and $m \geq 4$. □

By Observations 3.2 and Lemmas 3.3 and 3.4, we can induce a general expression for $n \geq 7$.

Lemma 3.5. *For $n \geq 7$,*

$$\begin{aligned}
&\nu(D_n) \\
&\leq \left(\tfrac{(n-1)!}{5!}\right)^2 \nu(D_6) \\
&\quad + \sum_{i=2j+1, j\geq 3}^{n} \left(\left(\tfrac{(n-1)!}{(i-1)!}\right)^2 \cdot \tfrac{(i-1)!}{24}(3i^4 - 25i^3 + 75i^2 - 95i + 42) \right) \\
&\quad + \sum_{i=2j, j\geq 4}^{n} \left(\left(\tfrac{(n-1)!}{(i-1)!}\right)^2 \cdot \tfrac{(i-2)!}{24}(3i^5 - 28i^4 + 103i^3 - 188i^2 + 164i - 48) \right).
\end{aligned}$$

Proof. By Observation 3.2 and Lemma 3.3, we have a recursive expression for $n = 2m - 1$ and $m \geq 4$

$$\begin{aligned}
&\nu(D_n) \\
&= 6\nu(D'_n) \\
&\leq 6\Big\{ (n-1)^2 \nu(D'_{n-1}) \\
&\quad + \tfrac{(n-1)!}{144}[3(n-1)^4 - 13(n-1)^3 + 18(n-1)^2 - 8(n-1)] \Big\} \\
&= (n-1)^2 \nu(D_{n-1}) + \tfrac{(n-1)!}{24}\left(3n^4 - 25n^3 + 75n^2 - 95n + 42\right) \\
&= (n-1)^2 \nu(D_{n-1}) + \sum_{i=n}^{n} \left(\left(\tfrac{(n-1)!}{(i-1)!}\right)^2 \cdot \tfrac{(i-1)!}{24}(3i^4 - 25i^3 + 75i^2 - 95i + 42) \right).
\end{aligned}$$

By Observation 3.2 and Lemma 3.4, we have a recursive expression for $n = 2m$ and $m \geq 4$

$$\begin{aligned}
&\nu(D_n) \\
&= 6\nu(D'_n) \\
&\leq 6\Big\{ (n-1)^2 \nu(D'_{n-1}) \\
&\quad + \tfrac{(n-2)!}{144}[3(n-1)^5 - 13(n-1)^4 + 21(n-1)^3 - 17(n-1)^2 + 6] \Big\} \\
&= (n-1)^2 \nu(D_{n-1}) + \tfrac{(n-2)!}{24}\left(3n^5 - 28n^4 + 103n^3 - 188n^2 + 164n - 48\right) \\
&= (n-1)^2 \nu(D_{n-1}) \\
&\quad + \sum_{i=n}^{n} \left(\left(\tfrac{(n-1)!}{(i-1)!}\right)^2 \cdot \tfrac{(i-2)!}{24}(3i^5 - 28i^4 + 103i^3 - 188i^2 + 164i - 48) \right).
\end{aligned}$$

Then we proceed by induction on n.

1) The base case is $n = 7$.

$$
\begin{aligned}
&\nu(D_7)\\
&\le 6^2\nu(D_6) + \sum_{i=7}^{7}\left(\left(\tfrac{(n-1)!}{(i-1)!}\right)^2 \cdot \tfrac{(i-1)!}{24}(3i^4 - 25i^3 + 75i^2 - 95i + 42)\right)\\
&= \left(\tfrac{(n-1)!}{5!}\right)^2 \nu(D_6)\\
&\quad + \sum_{i=2j+1, j\ge 3}^{n}\left(\left(\tfrac{(n-1)!}{(i-1)!}\right)^2 \cdot \tfrac{(i-1)!}{24}(3i^4 - 25i^3 + 75i^2 - 95i + 42)\right)\\
&\quad + \sum_{i=2j, j\ge 4}^{n}\left(\left(\tfrac{(n-1)!}{(i-1)!}\right)^2 \cdot \tfrac{(i-2)!}{24}(3i^5 - 28i^4 + 103i^3 - 188i^2 + 164i - 48)\right).
\end{aligned}
$$

2) We assume this lemma is true for $n = k - 1$.

For $n = k = 2m$ and $m \ge 4$, we have

$$
\begin{aligned}
&\nu(D_k)\\
&\le (k-1)^2\nu(D_{k-1})\\
&\quad + \sum_{i=k}^{k}\left(\left(\tfrac{(k-1)!}{(i-1)!}\right)^2 \cdot \tfrac{(i-2)!}{24}(3i^5 - 28i^4 + 103i^3 - 188i^2 + 164i - 48)\right)\\
&= (k-1)^2\left[\left(\tfrac{(k-2)!}{5!}\right)^2 \nu(D_6)\right.\\
&\quad + \sum_{i=2j+1, j\ge 3}^{k-1}\left(\left(\tfrac{(k-2)!}{(i-1)!}\right)^2 \cdot \tfrac{(i-1)!}{24}(3i^4 - 25i^3 + 75i^2 - 95i + 42)\right)\\
&\quad \left. + \sum_{i=2j, j\ge 4}^{k-1}\left(\left(\tfrac{(k-2)!}{(i-1)!}\right)^2 \cdot \tfrac{(i-2)!}{24}(3i^5 - 28i^4 + 103i^3 - 188i^2 + 164i - 48)\right)\right]\\
&\quad + \sum_{i=k}^{k}\left(\left(\tfrac{(k-1)!}{(i-1)!}\right)^2 \cdot \tfrac{(i-2)!}{24}(3i^5 - 28i^4 + 103i^3 - 188i^2 + 164i - 48)\right)\\
&= \left(\tfrac{(k-1)!}{5!}\right)^2 \nu(D_6)\\
&\quad + \sum_{i=2j+1, j\ge 3}^{k}\left(\left(\tfrac{(k-1)!}{(i-1)!}\right)^2 \cdot \tfrac{(i-1)!}{24}(3i^4 - 25i^3 + 75i^2 - 95i + 42)\right)\\
&\quad + \sum_{i=2j, j\ge 4}^{k}\left(\left(\tfrac{(k-1)!}{(i-1)!}\right)^2 \cdot \tfrac{(i-2)!}{24}(3i^5 - 28i^4 + 103i^3 - 188i^2 + 164i - 48)\right).
\end{aligned}
$$

For $n = k = 2m + 1$ and $m \ge 4$, we have

$$
\begin{aligned}
&\nu(D_k)\\
&\le (k-1)^2\nu(D_{k-1}) + \sum_{i=k}^{k}\left(\left(\tfrac{(k-1)!}{(i-1)!}\right)^2 \cdot \tfrac{(i-1)!}{24}(3i^4 - 25i^3 + 75i^2 - 95i + 42)\right)
\end{aligned}
$$

$$
\begin{aligned}
&= (k-1)^2 \left[\left(\tfrac{(k-2)!}{5!} \right)^2 \nu(D_6) \right. \\
&\quad + \sum_{i=2j+1, j\geq 3}^{k-1} \left(\left(\tfrac{(k-2)!}{(i-1)!} \right)^2 \cdot \tfrac{(i-1)!}{24} (3i^4 - 25i^3 + 75i^2 - 95i + 42) \right) \\
&\quad \left. + \sum_{i=2j, j\geq 4}^{k-1} \left(\left(\tfrac{(k-2)!}{(i-1)!} \right)^2 \cdot \tfrac{(i-2)!}{24} (3i^5 - 28i^4 + 103i^3 - 188i^2 + 164i - 48) \right) \right] \\
&\quad + \sum_{i=k}^{k} \left(\left(\tfrac{(k-1)!}{(i-1)!} \right)^2 \cdot \tfrac{(i-1)!}{24} (3i^4 - 25i^3 + 75i^2 - 95i + 42) \right) \\
&= \left(\tfrac{(k-1)!}{5!} \right)^2 \nu(D_6) \\
&\quad + \sum_{i=2j+1, j\geq 3}^{k} \left(\left(\tfrac{(k-1)!}{(i-1)!} \right)^2 \cdot \tfrac{(i-1)!}{24} (3i^4 - 25i^3 + 75i^2 - 95i + 42) \right) \\
&\quad + \sum_{i=2j, j\geq 4}^{k} \left(\left(\tfrac{(k-1)!}{(i-1)!} \right)^2 \cdot \tfrac{(i-2)!}{24} (3i^5 - 28i^4 + 103i^3 - 188i^2 + 164i - 48) \right).
\end{aligned}
$$

By 1) and 2), Lemma 3.5 holds for all $n \geq 7$. □

By Lemma 3.5, for $n \geq 7$ we have

$$
\begin{aligned}
&cr(B_n) \\
&\leq \nu(D_n) \\
&\leq \left(\tfrac{(n-1)!}{5!} \right)^2 \nu(D_6) \\
&\quad + \sum_{i=2j+1, j\geq 3}^{n} \left(\left(\tfrac{(n-1)!}{(i-1)!} \right)^2 \cdot \tfrac{(i-1)!}{24} (3i^4 - 25i^3 + 75i^2 - 95i + 42) \right) \\
&\quad + \sum_{i=2j, j\geq 4}^{n} \left(\left(\tfrac{(n-1)!}{(i-1)!} \right)^2 \cdot \tfrac{(i-2)!}{24} (3i^5 - 28i^4 + 103i^3 - 188i^2 + 164i - 48) \right) \\
&= \tfrac{433}{1200} ((n-1)!)^2 + \sum_{i=2j+1, j\geq 3}^{n} \left[\tfrac{((n-1)!)^2}{24(i-1)!} (3i^4 - 25i^3 + 75i^2 - 95i + 42) \right] \\
&\quad + \sum_{i=2j, j\geq 4}^{n} \left[\tfrac{((n-1)!)^2}{24(i-1)!(i-1)} (3i^5 - 28i^4 + 103i^3 - 188i^2 + 164i - 48) \right] \\
&= \tfrac{433}{1200} ((n-1)!)^2 + \sum_{i=2j+1, j\geq 3}^{n} \tfrac{((n-1)!)^2}{24} \cdot \left[\tfrac{3}{(i-5)!} + \tfrac{5}{(i-4)!} \right] \\
&\quad + \sum_{i=2j, j\geq 4}^{n} \tfrac{((n-1)!)^2}{24} \cdot \left[\tfrac{3}{(i-5)!} + \tfrac{5}{(i-4)!} + \tfrac{3}{(i-3)!} - \tfrac{6}{(i-2)!} + \tfrac{6}{(i-1)!(i-1)} \right] \\
&= \tfrac{433}{1200} ((n-1)!)^2 + \tfrac{((n-1)!)^2}{24} \cdot \left[\sum_{i=7}^{n} \left(\tfrac{3}{(i-5)!} + \tfrac{5}{(i-4)!} \right) \right. \\
&\quad \left. + \sum_{i=2j, j\geq 4}^{n} \left(\tfrac{3}{(i-3)!} - \tfrac{6}{(i-2)!} + \tfrac{6}{(i-1)!(i-1)} \right) \right]
\end{aligned}
$$

$$
\begin{aligned}
&= \tfrac{433}{1200}((n-1)!)^2 + \tfrac{((n-1)!)^2}{24} \cdot \left[\left(\tfrac{3}{2} + \sum_{i=3}^{n-5} \tfrac{8}{i!} + \tfrac{5}{(n-4)!} \right) \right. \\
&\quad \left. + \sum_{i=2j,j\geq 4}^{n} \left(\tfrac{3}{(i-3)!} - \tfrac{6}{(i-2)!} + \tfrac{6}{(i-1)!(i-1)} \right) \right] \\
&= \tfrac{127}{300}((n-1)!)^2 + \tfrac{((n-1)!)^2}{24} \cdot \left[\tfrac{5}{(n-4)!} + \sum_{i=3}^{n-5} \tfrac{8}{i!} \right. \\
&\quad \left. + \sum_{i=2j,j\geq 4}^{n} \left(\tfrac{3}{(i-3)!} - \tfrac{6}{(i-2)!} + \tfrac{6}{(i-1)!(i-1)} \right) \right] \\
&= ((n-1)!)^2 \left[\tfrac{127}{300} + \tfrac{5}{24(n-4)!} + \sum_{i=3}^{n-5} \tfrac{1}{3i!} \right. \\
&\quad \left. + \sum_{i=2j,j\geq 4}^{n} \left(\tfrac{1}{8(i-3)!} - \tfrac{1}{4(i-2)!} + \tfrac{1}{4(i-1)!(i-1)} \right) \right].
\end{aligned}
$$

Therefore, Theorem 3.2 holds for all $n \geq 7$.

Acknowledgements

The authors wish to express their appreciation to the referee for his or her valuable comments and suggestions, which have led to improvements in the presentation of this paper.

References

[1] Akers, S.B., Krishnamurthy, B., A group theoretic model for symmetric interconnection networks, *IEEE Transactions on Computers* 38 (4) (1989) 555–566.

[2] Araki, T., Kikuchi, Y., Hamiltonian laceability of bubble-sort graphs with edge faults, *Inform. Sci.* 177 (13) (2007) 2679–2691.

[3] Bhatt, S.N., Leighton, F.T., A framework for solving VLSI graph layout problems, *J. Comput. System Sci.* 28 (1984) 300–343.

[4] Cheng, E., Lipták, L., Sherman, D., Matching preclusion for the (n,k)-bubble-sort graphs, *Int. J. Comput. Math.* 87 (11) (2010) 2408–2418.

[5] Erdős, P., Guy, R.K., Crossing number problems, *Amer. Math. Monthly* 80 (1973) 52–58.

[6] Faria, L., Figueiredo, C.M.H. de, Sýkora, O., Vrt'o, I., An improved upper bound on the crossing number of the hypercube, *J. Graph Theory* 59 (2008) 145–159.

[7] Garey, M.R., Johnson, D.S., Crossing number is NP-complete, *SIAM J. Alg. Disc. Math.* 4 (1983) 312–316.

[8] Lin, X., Yang, Y., Lü, J., Hao, X., The crossing number of $C(n; \{1, \lfloor n/2 \rfloor - 1\})$. *Util. Math.* 71 (2006) 245–255.

[9] Shih, Y.-K., Lin, C.-K., Hsu, D.F., Tan, J.J.M., Hsu, L.-H., The construction of mutually independent Hamiltonian cycles in bubble-sort graphs, *Int. J. Comput. Math.* 87 (10) (2010) 2212–2225.

[10] Zheng, W., Lin, X., Yang, Y., Deng, C., The crossing number of Knödel graph $W_{3,n}$. *Util. Math.* 75 (2008) 211–224.